



Guaranteed Analysis

Total Nitrogen (N)	15.00%
10.0% Water Soluble Nitrogen	
5.0% Water Insoluble Nitrogen	
Soluble Potash (K ₂ O)	3.00%
Iron (Fe).....	1.00%
1.00% Water Soluble Iron	

Derived from Feather, Fish, Blood Meals,
Potassium Sulfate and Iron Sulfate.

Plant Food's 15-0-3 Fairway Grade provides a cost-effective fairway, lawncare and landscape natural based fertilizer product. This product combines the nitrogen benefits from feather and blood meals. This combination offers excellent residual color and growth that ranges from 6 to 10 weeks pending the applied rate.

15-0-3 Fairway Grade, 180 SGN

Directions for Use:

Fairway Maintenance Rate: Apply 150 - 250 lbs. per Acre for residual color ranging from 6 to 10 weeks. Apply 2 to 4 times per year.

Seasonal Lawncare Maintenance Rates:

Spring: Apply 5 lbs. per 1,000 sq. ft. for color, growth and rooting.

Summer: Apply 4 lbs. per 1,000 sq. ft. for stress and wear tolerance.

Fall: Apply 5.5 lbs. per 1,000 sq. ft. for rooting and spring color.

Watering Directions:

The product can be applied to wet or dry or stressed turf without any problems. For product breakdown, the key is WET – DRY – MOW. The product is applied and lightly watered – one cycle or revolution of the irrigation head. These products have a sugar binder. It is held together by sugars under pressure from the manufacturing process. The water aids in dissolving the sugar and releasing the pressure. The granular will begin to swell as the product dries. The product is ready to mow when dry to touch. It is recommended to mow without baskets to avoid picking up the product. For shorter mowing heights, a lower application rate at more frequent intervals is recommended. A dew whip can enhance breakdown and speed mowing.

Application Rates for 15-0-3, Fairway Grade, 180 SGN

Lbs/ 1,000 sq. ft.	Lbs/ One Acre	Bags/ One Acre	Nitrogen/ 1,000 sq. ft.	Potassium/ 1,000 sq. ft.	Iron/ 1,000 sq. ft.	Spreader Settings			
						Scotts R8	Lesco	Lely	Vicon
3.4 lb.	146 lbs.	2.93 bags	0.50	0.10	0.0336	I	E	3	30
6.7 lb.	290 lbs.	5.79 bags	1.00	0.20	0.0665	R	I	6	49

Spreader settings are an initial guideline for calibrating the spreader.

Caution:

Keep away from children and domestic animals. Avoid contact with eyes, open cuts, or sores. Harmful if swallowed. External: Flood with water. Internal: Induce vomiting. Contact a physician immediately.

Storage and Handling:

Store in a cool, dry place. Keep container tightly closed. Do not add water or other material to the container. Do not contaminate water, food, or feed by storage or disposal. Do not store near acids or other acidic materials.

Available Container Sizes:

50 lb. Bag (22.7 kg)

