

**PLANT SAFETY/COMPATIBILITY:**

**K-PHITE 7LP Fungicide** is compatible with most pesticide products used in turf and ornamentals. **K-PHITE 7LP Fungicide** shows no phytotoxicity on plants when used as instructed on the label. Always conduct a phytotoxicity check before applying to large areas. **K-PHITE 7LP Fungicide has a 4 hour REI and a 0 day pre-harvest.**

**GENERAL APPLICATION INSTRUCTIONS:**

**Chemigation:** Apply K-PHITE 7LP Fungicide through microjet, drip, solid set and centerpivot sprinkler irrigation systems. Apply **K-PHITE 7LP Fungicide** by various application methods, including foliar sprays, trunk injection, soil drench, soil incorporation and bare root dip. For foliar sprays, apply **K-PHITE 7LP Fungicide** with sufficient water for adequate coverage of foliage, according to crop and growth stage. To insure good coverage, spray wet.

**INSTRUCTIONS FOR APPLICATIONS****ORNAMENTALS, BEDDING PLANTS, TURF and TREES (CONIFERS)  
HOME AND GARDEN**

**K-PHITE 7LP Fungicide** is used for control or prevention of Bacterial Blight, Bacterial Leafspot, Pythium, Downy Mildew, Powdery Mildew, Erwinia, Fusarium, Phytophthora and various other diseases grown in commercial nurseries and greenhouses, golf courses, parks and commercial landscapes. K-PHITE 7LP Fungicide may be applied to all ornamental and landscape plants including (but not limited to): Aglaonema, Aphelandra, Arbovitae, Ash, Azaleas, Bougainvilleas, Boxwood, Bushes, Cattleya Skinneri, Cedars, Ceanothus, Cotoneaster, Cissus, Diffenbachia, Dogwood, Ferns, Ficus, Flowering Trees, Hibiscus, Hard Wood Trees, Holly, Home Lawns, Juniper, Magnolia Trees, Maple, Pines, Oaks, Palms, Peperomia, Photinia, Pittosporum, Philodendron, Pieris, Pothos, Red woods, Rhododendron, Roses, Schefflera, Sedum, Sempervivum, Shrubs, Spathiphyllum, Sycamore, Taxus media, Zygocactus.

**BEDDING PLANTS:** Ageratum, Algerian Ivy, Anthurium, Artemisia, Aster, Baby's Breath, Begonia, Caladium, Carnation, Chrysanthemum, Coleus, Columbine, Daisy, Daylily, Delphinium, Easter Lily, English Ivy, Foxglove, Gaillardia, Geranium, Gloxinia, Impatiens, Marigold, Pansy, Petunia, Phlox, Pinks, Poinsettia, Primrose, Prostrate Rosemary, Salvia, Snapdragon, Verbena, Vinca, Zinnia

**TREES:** Douglas Fir, Spruce, Pines and Oaks.

**TURF:** **K-PHITE 7LP Fungicide** is a systemic fungicide, which is used for the control of Pythium blight, Summer Complex (northern and southern turf types), Dollar Spot, Grey Leaf Spot, Brown Patch, Anthracnose and damping-off diseases to turf grasses on sod farms, commercial turf production, golf courses, parks and commercial landscapes, home lawns.

Applications should be made prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe.

Apply **K-PHITE 7LP Fungicide** prior to disease development in conjunction with good cultural practices. Use the higher rate when disease pressure is severe. Do not exceed recommended rates or apply more frequently than the specified intervals to avoid plant injury. Do not apply to plants that are dormant or under heat or moisture stress.

Crops	Disease	Application
<b>Ornamentals</b> Bedding Plants Flowers (Greenhouse and Field), Greenhouse Ornamentals Greenhouse Vegetables, Bare Root Stock, Container Stock Fresh Cut Flowers	<b>Phytophthora</b> (Root Rot)	Foliar 1 to 2 qts./100 gal. of water or 1/3 fl. oz. to 2/3 fl. oz. per gal. of water. (2.5 to 5 L/1000 L of water)
	<b>Fusarium</b>	
	<b>Pythium</b> (Damping Off)	
	<b>Rhizoctonia</b> (Crown Root)	
	<b>Erwinia</b> (Fire Blight)	Soil Drench 12-24oz/100 gal. of water
	<b>Xanthomonas</b> (Bacterial Blight) (Bacteria Leaf Spot)	
	<b>Downy Mildew</b>	Foliar Spray 3-6 fl. oz./1,000 sq. ft.
	<b>Powdery Mildew</b>	or
	<b>Ralstonia spp.</b> (Geranium Wilt)	3 to 6 qts./100 gal. of water/acre.
	<b>Pythium</b> (Yellow Turf)	Irrigation Overhead: 2-3 qts./acre. (5-7 L/ Ha) in at least 100 gallons (1000 L) of water.
<b>Turf</b>	<b>Rhizoctonia</b> (Brown Patch)	
	<b>Sclerotinia</b> (Dollar Spot)	
	<b>Helminthosporium</b> (Gray Leaf Spot)	
	<b>Alternaria</b>	
	<b>Anthrachnose</b>	
	<b>Phytophthora</b> (Sudden Oak Death) (Root Rot)	Foliar Spray 1 to 2 qts./100 gal. water /acre (2.5 to 5 L/1000 L water/Ha) or 1/3 fl. oz. to 2/3 fl. oz. per gal. Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed.
		Trunk Injections 3 teaspoons /linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH) (15 milliliters/ linear meter of canopy width or 5 centimeters DBH) Inject directly into tree.
<b>Conifers and Trees</b>		

**NOTICE TO BUYER:**

To the extent permitted by law, all conditions and warranties, and statutory or other rights of action which buyer or any other user may have against Plant Food Systems or Seller are hereby excluded. Plant Food Systems hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information provided by Plant Food Systems or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Plant Food Systems' liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price thereof.

**Plant Food Systems**

P.O. Box 775  
Zellwood, Florida 32798  
407-889-7755  
www.plantfoodsystems.com

Always read and follow label directions.  
Specimen label only, read and follow registered label directions.

**K-PHITE 7LP**  
SYSTEMIC FUNGICIDE BACTERICIDE

Safe, dependable crop protection.